



HAL
open science

Influence of nitrogen fertilisation and autumn mowing on dual use of thinopyrum intermedium

Fagnant Laura, Olivier Duchene, David Christophe, Bodson Bernard, Beckers
Yves, Bindelle Jérôme, Dumont Benjamin

► **To cite this version:**

Fagnant Laura, Olivier Duchene, David Christophe, Bodson Bernard, Beckers Yves, et al.. Influence of nitrogen fertilisation and autumn mowing on dual use of thinopyrum intermedium. ASA, CSSA and SSSA International Annual Meetings, 2020, Phoenix, Arizona, United States. hal-03654385

HAL Id: hal-03654385

<https://isara.hal.science/hal-03654385v1>

Submitted on 28 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INFLUENCE OF NITROGEN FERTILISATION AND AUTUMN MOWING ON DUAL USE OF *THINOPYRUM INTERMEDIUM*.

Fagnant Laura¹, Duchêne Olivier², Roiseux Olivier³, David Christophe², Bodson Bernard¹, Beckers Yves¹, Bindelle Jérôme¹, Dumont Benjamin¹.

¹ Liege University - Gembloux Agro-Bio Tech, TERRA Res. and Teach. center, Plant Sciences, Crop Sciences, Gembloux, Belgium.

Laura.fagnant@uliege.be

² ISARA, Agroecology and Environment Research Unit, Lyon, France.

³ Walagri, Sombrefe, Belgium.

The double production grain-forage of *Thinopyrum intermedium* (Host) Barkworth & D.R. Dewey for several years could be of major interest for crop diversification and environmental improvement of the highly productive agroecosystems in Belgium. Indeed, the perennality of the species could be a way to combine sustainability and agricultural productions. Grain production of *Th. intermedium* is limited compared to annual cereals, but can be offset by its above-ground biomass production used as forage. As the crop has never been implanted in our regions, characterizing its potential is crucial. The aim of this research is to optimize the dual use of the crop by studying different agronomic levers such as nitrogen fertilisation and autumn mowing. Our results show that the nitrogen fertilisation is very useful in the establishment year, but does not seem significant the subsequent years. During first year, N fertilization has increased grain yield from 0,8 to 1,5 tons/ha and aerial biomass production from 4,3 to 10,6 tons/ha of DM with a fertilisation of 100 kgN/ha applied before stem elongation (BBCH 30). The aestival grain productions can furthermore be combined with post-harvest biomass regrowth that can be mowed in autumn. This regrowth represented roughly 1,3tons/ha of DM during the first year. Although decreasing the amount of total above-ground biomass the following season, this mowing allows to valorise an aerial biomass of good feed values. This cut could also have a positive effect when the nitrogen fertilisation is high by reducing intraspecific competition. These results must be confirmed in the coming years, but they constitute a first understanding of field management to optimize dual production which must be coordinated with grain production objectives.

Keywords : *Thinopyrum intermedium*, perennial grains, nitrogen, mowing