

Combining wheat and prairies using functional diversity to promote positive ecological interactions and a better regulation of C and N cycles in agroecosystems

Thomas Bécu, Elodie Pascon, Sébastien Fontaine, Audrey Niboyet, Gaël Alvarez, Sébastien Barot, Lisa Besson, Olivier Duchene, Amélie Cantarel, Xavier Le Roux, et al.

▶ To cite this version:

Thomas Bécu, Elodie Pascon, Sébastien Fontaine, Audrey Niboyet, Gaël Alvarez, et al.. Combining wheat and prairies using functional diversity to promote positive ecological interactions and a better regulation of C and N cycles in agroecosystems. SFE2 Congress, Oct 2024, Lyon, France. 2024. hal-04804258

HAL Id: hal-04804258 https://isara.hal.science/hal-04804258v1

Submitted on 26 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Combining wheat and prairies using functional diversity to promote positive ecological interactions and a better regulation of C and N cycles in agroecosystems

Thomas Bécu¹, Elodie Pascon¹, Sébastien Fontaine¹, Audrey Niboyet^{2,3}, Gael Alvarez¹, Sebastien Barot², Lisa Besson¹, Amélie Cantarel⁴, Olivier Duchene⁵, Xavier Le Roux⁴, Nicolas Gross¹

- ¹ INRAE, VetAgro Sup, Université Clermont Auvergne, UMR Ecosystème Prairial (UREP), Clermont-Ferrand 63000, France
- ² Sorbonne Université, Université Paris Cité, UPEC, CNRS, INRAE, IRD, Institut d'Ecologie et des Sciences de l'Environnement de Paris (IEES-Paris), Paris 75005, France

³ AgroParisTech, Palaiseau 91120, France


⁴ INRAE, CNRS, Université Lyon 1, Laboratoire d'Ecologie Microbienne (LEM), UMR 1418 INRAE, Villeurbanne 69622, France


⁵ ISARA, Agroecology and Environment Research Unit, Lyon 69364, France

Context


Plant functional diversity interacts with soil microbial communities, resulting in a heterarchical network that drives soil biogeochemical cycles ^{1–3}. In permanent and diverse temperate prairies, these interactions generate a tight coupling between soil carbon (C) and nitrogen (N) cycles and a coordination between plant nutrient demand and soil offer, that favours the resource use efficiency 4. This emerging property could be used in highly diversified agroecosystems to increase the biomass production of a target species while improving carbon storage and nutrient recycling. To explore this opportunity, we combined an annual crop (winter wheat) with permanent and diverse prairies cultivated in alternating strips, in new agricultural systems called "agroprairies". However, diversification also implies plant-plant interactions that are driven by plant functional traits as well as by management practices, and that will affect the final performance of a target species 5,6.


How to promote positive plant-plant interactions and reduce competition in highly diverse agroecosystems, with the help of functional traits distribution?














What is the effect of prairie functional structure, fertilisation management and wheat variety on wheat nutrition at the onset of stem elongation?


Main Results

- Variable effect of prairies was observed on wheat N_{up} (N in upper leaves): 43 plots with a positive effect, 117 plots with a negative effect
- Wheat response to prairies was highly impacted by wheat variety and its interaction with fertilisation (23.2 and 22.5% of variance explained), with a significant effect of prairie functional characteristics
- We observed an overall positive effect of legume abundance and fast growing strategy, and a negative effect of trait dispersion (LA, LDMC, H) on wheat response to prairies
- The positive effect of fast growing strategy on the N_{up} response in Wheat was lowered by fertilisation and differed between wheat varieties.









Conclusion

Our results showed a variable and significant effect of prairie functional structure on the response of wheat nutrition at the onset of stem elongation, depending on N management. Together with forthcoming data on height and biomass production, our study provides first insights into the potential benefits of combining an annual crop with diverse prairies, along with key factors that will help to design agroecosystems that maximise the benefits of functional diversity while minimising negative interactions.









References

Abalos et. al, 2019, Journal of Applied Ecology De Deyn et. al, 2008, Ecology Letters Fernandez et. al, 2022, Journal of Ecology

Gross et. al, 2007, Journal of Ecology

7. Le Bagousse-Pinguet et. al, 2021, PNAS

8. Gross et. al, 2021, Ecology Letters

- Fontaine et. al, 2024, Global Change Biology Holden et. al, 2024, Functional Ecology